

1230372

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

September 24, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/498,309

FILING DATE: August 28, 2003

RELATED PCT APPLICATION NUMBER: PCT/US04/26910

Certified by



Jon W Dudas

Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office



Best Available Copy

18334 U.S. PTO
09/28/03

PTO/SB/16 (08-03)
Approved for use through 07/31/2006. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No.

17497 U.S. PTO
06/28/03

INVENTOR(S)					
Given Name (first and middle (if any))		Family Name or Surname		Residence (City and either State or Foreign Country)	
David S. L. Gordon		Garvey Letts		Dover, MA Dover, MA	
Additional inventors are being named on the <u>1</u> separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
Nitrosated and/or Nitrosylated Cardiovascular Compounds, Compositions and Methods of Use					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input checked="" type="checkbox"/> Customer Number: 25270					
OR					
<input type="checkbox"/> Firm or Individual Name		Edward D. Grieff			
Address		Hale and Dorr LLP			
Address		1455 Pennsylvania Avenue, NW			
City		Washington	State	DC	Zip 20004
Country		US	Telephone	Fax	2029428484
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages <u>45</u>		<input type="checkbox"/> CD(s), Number _____			
<input type="checkbox"/> Drawing(s) Number of Sheets _____		<input checked="" type="checkbox"/> Other (specify) <u>Return Postcard</u>			
<input type="checkbox"/> Application Date Sheet. See 37 CFR 1.76					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. <input type="checkbox"/> A check or money order is enclosed to cover the filing fees. <input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: <u>08-0219</u> <input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.				FILING FEE Amount (\$) <div style="border: 1px solid black; padding: 10px; text-align: center;">\$80.00</div>	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government. <input checked="" type="checkbox"/> No. <input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

Respectfully submitted

SIGNATURE

TYPED or PRINTED NAME Edward D. Grieff

TELEPHONE 202-942-8400

Date August 28, 2003

REGISTRATION NO. 38,898

(if appropriate)

Docket Number: 102258

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

PROVISIONAL APPLICATION COVER SHEET
Additional Page

PTO/SB/16 (02-01)

Approved for use through 10/31/2002. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Docket Number

102258.

INVENTOR(S)/APPLICANT(S)

Given Name (first and middle [if any])	Family or Surname	Residence (City and either State or Foreign Country)
Manuel	Worcel	Boston, MA

Number 2 of 2

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

NITROSATED AND/OR NITROSYLATED CARDIOVASCULAR COMPOUNDS, COMPOSITIONS AND METHODS OF USE

FIELD OF THE INVENTION

The invention describes novel nitrosated and/or nitrosylated cardiovascular compounds
5 and pharmaceutically acceptable salts thereof, and novel compositions comprising at least one
nitrosated and/or nitrosylated cardiovascular compound, and, optionally, at least one compound
that donates, transfers or releases nitric oxide, stimulates endogenous synthesis of nitric oxide,
elevates endogenous levels of endothelium-derived relaxing factor or is a substrate for nitric
oxide synthase, and/or at least one therapeutic agent. The invention also provides novel
10 compositions comprising at least one cardiovascular compound, and at least one compound that
donates, transfers or releases nitric oxide, elevates endogenous levels of endothelium-derived
relaxing factor, stimulates endogenous synthesis of nitric oxide or is a substrate for nitric oxide
synthase and/or at least one therapeutic agent. The invention also provides novel kits comprising
at least one cardiovascular compound, that is optionally nitrosated and/or nitrosylated, and,
15 optionally, at least one nitric oxide donor and/or at least one therapeutic agent. The invention
also provides methods for treating cardiovascular diseases, methods for treating diseases
resulting from oxidative stress and methods for treating endothelial dysfunctions. The nitrosated
and/or nitrosylated cardiovascular compounds are preferably nitrosated and/or nitrosylated
aldosterone antagonists, nitrosated and/or nitrosylated angiotensin II antagonists, nitrosated
20 and/or nitrosylated hydralazine compounds, nitrosated and/or nitrosylated calcium channel
blockers, nitrosated and/or nitrosylated diuretics, nitrosated and/or nitrosylated endothelin
antagonists, nitrosated and/or nitrosylated neutral endopeptidase inhibitors and nitrosated and/or
nitrosylated renin inhibitors.

BACKGROUND OF THE INVENTION

25 The decline in cardiovascular morbidity and mortality in the United States over the past
three decades has been the result of significant advances in research on cardiovascular disease
mechanisms and therapeutic strategies. The incidence and prevalence of myocardial infarction
and death from myocardial infarction, as well as that from cerebrovascular accident, have
decreased significantly over this period largely owing to advances in prevention, early diagnosis,
30 and treatment of these very common diseases.

The compounds administered for the treatment of cardiovascular diseases, diseases

resulting from oxidative and/or endothelial dysfunctions often result in toxicity, chronic and/or debilitating side effects. Cardiovascular compounds such as ACE inhibitors, beta-adrenergic blockers, antithrombotic and vasodilator drugs or anti-hyperlipidemic drugs, show, for example, respiratory toxicity resulting in asthma and/or bronchitis. Hence there is a need in the art for cardiovascular compounds that have improved efficacy, lower toxicity and that can be used at low dosages. The invention is directed to these, as well as other, important ends.

SUMMARY OF THE INVENTION

The invention provides novel cardiovascular compounds that are substituted with at least one NO and/or NO₂ group (i.e., nitrosylated and/or nitrosated), and pharmaceutically acceptable salts thereof. The cardiovascular compounds can be, for example, aldosterone antagonists, angiotensin II antagonists, hydralazine compounds, calcium channel blockers, diuretics, endothelin antagonists, neutral endopeptidase inhibitors and renin inhibitors. The cardiovascular compounds can be nitrosated and/or nitrosylated through one or more sites such as oxygen (hydroxyl condensation), sulfur (sulfhydryl condensation) and/or nitrogen. The invention also provides compositions comprising the novel compounds described herein in a pharmaceutically acceptable carrier.

The invention is also based on the discovery that administering at least one cardiovascular compound or pharmaceutically acceptable salts thereof, that is optionally substituted with at least one NO and/or NO₂ group (i.e., nitrosylated and/or nitrosated), and, optionally, at least one nitric oxide donor improves the properties of the cardiovascular compound. Nitric oxide donors include, for example, S-nitrosothiols, nitrites, nitrates, N-oxo-N-nitrosamines, SPM 3672, SPM 5185, SPM 5186 and analogues thereof, and substrates of the various isozymes of nitric oxide synthase. Thus, another embodiment of the invention provides compositions comprising at least one cardiovascular compound that is optionally substituted with at least one NO and/or NO₂ group (i.e., nitrosylated and/or nitrosated), and at least one compound that donates, transfers or releases nitric oxide as a charged species, i.e., nitrosonium (NO⁺) or nitroxyl (NO⁻), or as the neutral species, nitric oxide (NO•), and/or stimulates endogenous production of nitric oxide or EDRF *in vivo* and/or is a substrate for nitric oxide synthase. The invention also provides for such compositions in a pharmaceutically acceptable carrier.

Yet another embodiment of the invention provides compositions comprising at least one cardiovascular compound, that is optionally substituted with at least one NO and/or NO₂ group

(i.e., nitrosylated and/or nitrosated), and, optionally, at least one compound that donates, transfers or releases nitric oxide as a charged species, i.e., nitrosonium (NO^+) or nitroxyl (NO^-), or as the neutral species, nitric oxide ($\text{NO}\bullet$), and/or is a substrate for nitric oxide synthase and/or at least one therapeutic agent, including, but not limited to, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H_2 receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and mixtures of two or more thereof. The invention also provides for such compositions in a pharmaceutically acceptable carrier.

Yet another embodiment of the invention provides methods for treating cardiovascular diseases, methods for treating diseases resulting from oxidative stress and/or methods for treating endothelial dysfunctions in a patient in need thereof comprising administering to the patient a therapeutically effective amount of at least one cardiovascular compound, that is optionally substituted with at least one NO and/or NO_2 group (i.e., nitrosylated and/or nitrosated), and, optionally, at least one compound that donates, transfers or releases nitric oxide as a charged species, i.e., nitrosonium (NO^+) or nitroxyl (NO^-), or as the neutral species, nitric oxide ($\text{NO}\bullet$), and/or stimulates endogenous production of nitric oxide or EDRF *in vivo* and/or is a substrate for nitric oxide synthase (i.e. NO donor). The methods can optionally further comprise the administration of at least one therapeutic agent, such as, for example, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H_2 receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and mixtures of two or more thereof. In this embodiment of the invention, the methods can involve administering the nitrosated and/or nitrosylated cardiovascular compounds, administering the cardiovascular compounds, that are optionally nitrosated and/or nitrosylated,

and NO donors, administering the cardiovascular compounds, that are optionally nitrosated and/or nitrosylated, and therapeutic agents, or administering the cardiovascular compounds, that are optionally nitrosated and/or nitrosylated, NO donors, and therapeutic agents. The cardiovascular compounds, nitric oxide donors, and/or therapeutic agents can be administered separately or as components of the same composition in one or more pharmaceutically acceptable carriers.

In yet another embodiment the invention provides kits comprising at least one cardiovascular compound, that is optionally nitrosated and/or nitrosylated, and, optionally, at least one compound that donates, transfers or releases nitric oxide as a charged species, i.e., nitrosonium (NO^+) or nitroxyl (NO^-), or as the neutral species, nitric oxide ($\text{NO}\bullet$), and/or stimulates endogenous production of nitric oxide or EDRF *in vivo* and/or is a substrate for nitric oxide synthase. The kit can further comprise at least one therapeutic agent, such as, for example, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H_2 receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and mixtures of two or more thereof. The cardiovascular compound, the nitric oxide donor and/or therapeutic agent, can be separate components in the kit or can be in the form of a composition in one or more pharmaceutically acceptable carriers.

These and other aspects of the invention are described in detail herein.

DETAILED DESCRIPTION OF THE INVENTION

As used throughout the disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings.

“Cardiovascular diseases” refers to any cardiovascular disease, including but not limited to, congestive heart failure, hypertension (e.g. pulmonary hypertension, low-renin hypertension, salt-sensitive hypertension, low-renin, salt-sensitive hypertension, thromboembolic pulmonary hypertension; pregnancy-induced hypertension; renovascular hypertension; hypertension-dependent end-stage renal disease), left ventricular hypertrophy with disproportionate microvascularization, hypertrophic cardiomyopathy, diastolic dysfunction, myocardial

infarctions, cerebral infarctions, atherosclerosis, atherogenesis, thrombosis, ischemic heart disease, coronary artery diseases, acute or chronic renal failure, stable, unstable and variant (Prinzmetal) angina, hypercholesterolemia, cardiac edema, renal insufficiency, nephrotic edema, hepatic edema, stroke, transient ischemic attacks, cerebrovascular incidents, restenosis, platelet
5 adhesion, platelet aggregation, smooth muscle cell proliferation, supraventricular and ventricular arrhythmias, atrial fibrillation or atrial flutter, angina pectoris, diabetes secondary aldosteronism, primary and secondary pulmonary hyperaldosteronism, glomerulonephritis, scleroderma, glomerular sclerosis, proteinuria of primary renal disease, diabetic retinopathy, vascular complications associated with the use of medical devices, wounds associated with the use of
10 medical devices, pulmonary thromboembolism, cerebral thromboembolism, thrombophlebitis, thrombocytopenia, bleeding disorders, thrombotic or thromboembolic events, such as, for example, arterial thrombosis, coronary thrombosis, heart valve thrombosis, coronary stenosis, stent thrombosis and graft thrombosis and the like.

“Diseases resulting from oxidative stress” refers to any disease that involves the
15 generation of free radicals or radical compounds, such as, for example, atherogenesis, atheromatosis, arteriosclerosis, atherosclerosis, vascular hypertrophy associated with hypertension, hyperlipoproteinaemia, normal vascular degeneration through aging, parathyroidal reactive hyperplasia, chronic renal disease, neoplastic diseases, inflammatory diseases, neurological and acute bronchopulmonary disease, tumorigenesis, ischemia-reperfusion
20 syndrome, arthritis, sepsis, cognitive dysfunction, endotoxic shock, endotoxin-induced organ failure, and the like.

“Endothelial dysfunction” refers to the dysfunction of a patient’s endothelium that results in, for example, arteriosclerosis, congestive heart failure, hypertension, cardiovascular diseases, cerebrovascular diseases, renovascular diseases, mesenteric vascular diseases, pulmonary
25 vascular diseases, ocular vascular diseases, peripheral vascular diseases, and the like.

“Antioxidant” refers to and includes any compound that can react and quench a free radical.

“Angiotensin converting enzyme (ACE) inhibitor” refers to compounds that inhibit an enzyme which catalyzes the conversion of angiotensin I to angiotensin II. ACE inhibitors
30 include, but are not limited to, amino acids and derivatives thereof, peptides, including di- and tri-peptides, and antibodies to ACE which intervene in the renin-angiotensin system by inhibiting

the activity of ACE thereby reducing or eliminating the formation of the pressor substance angiotensin II.

“Angiotensin II antagonists” refers to compounds which interfere with the function, synthesis or catabolism of angiotensin II. Angiotensin II antagonists include peptide compounds and non-peptide compounds, including, but not limited to, angiotensin II antagonists, angiotensin II receptor antagonists, agents that activate the catabolism of angiotensin II, and agents that prevent the synthesis of angiotensin I from angiotensin II. The renin-angiotensin system is involved in the regulation of hemodynamics and water and electrolyte balance. Factors that lower blood volume, renal perfusion pressure, or the concentration of sodium in plasma tend to activate the system, while factors that increase these parameters tend to suppress its function.

“Anti-hyperlipidemic drugs” refers to any compound or agent that has the effect of beneficially modifying serum cholesterol levels such as, for example, lowering serum low density lipoprotein (LDL) cholesterol levels, or inhibiting oxidation of LDL cholesterol, whereas high density lipoprotein (HDL) serum cholesterol levels may be lowered, remain the same, or be increased. Preferably, the anti-hyperlipidemic drug brings the serum levels of LDL cholesterol and HDL cholesterol (and, more preferably, triglyceride levels) to normal or nearly normal levels.

“Neutral endopeptidase inhibitors” refers to and includes compounds that are antagonists of the renin angiotensin aldosterone system including compounds that are dual inhibitors of neutral endopeptidases and angiotensin converting (ACE) enzymes.

“Renin inhibitors” refers to compounds which interfere with the activity of renin.

“Platelet reducing agents” refers to compounds that prevent the formation of a blood thrombus via any number of potential mechanisms. Platelet reducing agents include, but are not limited to, fibrinolytic agents, anti-coagulant agents and any inhibitors of platelet function. Inhibitors of platelet function include agents that impair the ability of mature platelets to perform their normal physiological roles (i.e., their normal function, such as, for example, adhesion to cellular and non-cellular entities, aggregation, release of factors such as growth factors) and the like.

“Proton pump inhibitor” refers to any compound that reversibly or irreversibly blocks gastric acid secretion by inhibiting the H^+/K^+ -ATP ase enzyme system at the secretory surface of the gastric parietal cell.


"NSAID" refers to a nonsteroidal anti-inflammatory compound or a nonsteroidal anti-inflammatory drug. NSAIDs inhibit cyclooxygenase, the enzyme responsible for the biosyntheses of the prostaglandins and certain autocoid inhibitors, including inhibitors of the various isozymes of cyclooxygenase (including but not limited to cyclooxygenase-1 and -2), and as inhibitors of both cyclooxygenase and lipoyxygenase.

"Therapeutic agent" includes any therapeutic agent that can be used to treat or prevent the diseases described herein. "Therapeutic agents" include, for example, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H₂ receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitor, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and the like. Therapeutic agent includes the pro-drugs and pharmaceutical derivatives thereof including, but not limited to, the corresponding nitrosated and/or nitrosylated derivatives. Although nitric oxide donors have therapeutic activity, the term "therapeutic agent" does not include the nitric oxide donors described herein, since nitric oxide donors are separately defined.

"Cyclooxygenase-2 (COX-2) selective inhibitor" refers to a compound that selectively inhibits the cyclooxygenase-2 enzyme over the cyclooxygenase-1 enzyme. In one embodiment, the compound has a cyclooxygenase-2 IC₅₀ of less than about 2 μM and a cyclooxygenase-1 IC₅₀ of greater than about 5 μM, in the human whole blood COX-2 assay (as described in Brideau et al., *Inflamm Res.*, 45: 68-74 (1996)) and also has a selectivity ratio of cyclooxygenase-2 inhibition over cyclooxygenase-1 inhibition of at least 10, and preferably of at least 40. In another embodiment, the compound has a cyclooxygenase-1 IC₅₀ of greater than about 1 μM, and preferably of greater than 20 μM. The compound can also inhibit the enzyme, lipoyxygenase. Such selectivity may indicate an ability to reduce the incidence of common NSAID-induced side effects.

"Hydrazino" refers to H₂N-N(H)-.

"Hydralazine compound" refers to a compound having the formula:



wherein a, b and c are independently a single or double bond; R₁ and R₂ are each independently a hydrogen, an alkyl, an ester or a heterocyclic ring, wherein alkyl, ester and heterocyclic ring are as defined herein; R₃ and R₄ are each independently a lone pair of electrons or a hydrogen, with the proviso that at least one of R₁, R₂, R₃ and R₄ is not a hydrogen.

5 Exemplary hydralazine compounds include budralazine, cadralazine, dihydralazine, endralazine, hydralazine, pildralazine, todralazine, and the like.

"Patient" refers to animals, preferably mammals, most preferably humans, and includes males and females, and children and adults.

10 "Therapeutically effective amount" refers to the amount of the compound and/or composition that is effective to achieve its intended purpose.

"Transdermal" refers to the delivery of a compound by passage through the skin and into the blood stream.

"Transmucosal" refers to delivery of a compound by passage of the compound through the mucosal tissue and into the blood stream.

15 "Penetration enhancement" or "permeation enhancement" refers to an increase in the permeability of the skin or mucosal tissue to a selected pharmacologically active compound such that the rate at which the compound permeates through the skin or mucosal tissue is increased.

20 "Carriers" or "vehicles" refers to carrier materials suitable for compound administration and include any such material known in the art such as, for example, any liquid, gel, solvent, liquid diluent, solubilizer, or the like, which is non-toxic and which does not interact with any components of the composition in a deleterious manner.

25 "Sustained release" refers to the release of a therapeutically active compound and/or composition such that the blood levels of the therapeutically active compound are maintained within a desirable therapeutic range over an extended period of time. The sustained release formulation can be prepared using any conventional method known to one skilled in the art to obtain the desired release characteristics.

"Nitric oxide adduct" or "NO adduct" refers to compounds and functional groups which, under physiological conditions, can donate, release and/or directly or indirectly transfer any of the three redox forms of nitrogen monoxide (NO⁺, NO⁻, NO•), such that the biological activity of

the nitrogen monoxide species is expressed at the intended site of action.

"Nitric oxide releasing" or "nitric oxide donating" refers to methods of donating, releasing and/or directly or indirectly transferring any of the three redox forms of nitrogen monoxide (NO^+ , NO^- , NO^\bullet), such that the biological activity of the nitrogen monoxide species is expressed at the intended site of action.

"Nitric oxide donor" or "NO donor" refers to compounds that donate, release and/or directly or indirectly transfer a nitrogen monoxide species, and/or stimulate the endogenous production of nitric oxide or endothelium-derived relaxing factor (EDRF) *in vivo* and/or elevate endogenous levels of nitric oxide or EDRF *in vivo*. "NO donor" also includes compounds that are substrates for nitric oxide synthase.

"Alkyl" refers to a lower alkyl group, a haloalkyl group, a hydroxyalkyl group, an alkenyl group, an alkynyl group, a bridged cycloalkyl group, a cycloalkyl group or a heterocyclic ring, as defined herein. An alkyl group may also comprise one or more radical species, such as, for example a cycloalkylalkyl group or a heterocyclicalkyl group.

"Lower alkyl" refers to branched or straight chain acyclic alkyl group comprising one to about ten carbon atoms (preferably one to about eight carbon atoms, more preferably one to about six carbon atoms). Exemplary lower alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, pentyl, neopentyl, iso-amyl, hexyl, octyl, and the like.

"Substituted lower alkyl" refers to a lower alkyl group, as defined herein, wherein one or more of the hydrogen atoms have been replaced with one or more R^{100} groups, wherein each R^{100} is independently a hydroxy, an oxo, a carboxyl, a carboxamido, a halo, a cyano or an amino group, as defined herein.

"Haloalkyl" refers to a lower alkyl group, an alkenyl group, an alkynyl group, a bridged cycloalkyl group, a cycloalkyl group or a heterocyclic ring, as defined herein, to which is appended one or more halogens, as defined herein. Exemplary haloalkyl groups include trifluoromethyl, chloromethyl, 2-bromobutyl, 1-bromo-2-chloro-pentyl, and the like.

"Alkenyl" refers to a branched or straight chain $\text{C}_2\text{-C}_{10}$ hydrocarbon (preferably a $\text{C}_2\text{-C}_8$ hydrocarbon, more preferably a $\text{C}_2\text{-C}_6$ hydrocarbon) that can comprise one or more carbon-carbon double bonds. Exemplary alkenyl groups include propylenyl, buten-1-yl, isobutenyl, penten-1-yl, 2,2-methylbuten-1-yl, 3-methylbuten-1-yl, hexan-1-yl, hepten-1-yl, octen-1-yl, and

the like.

"Lower alkenyl" refers to a branched or straight chain C₂-C₄ hydrocarbon that can comprise one or two carbon-carbon double bonds.

"Substituted alkenyl" refers to a branched or straight chain C₂-C₁₀ hydrocarbon (preferably a C₂-C₈ hydrocarbon, more preferably a C₂-C₆ hydrocarbon) which can comprise one or more carbon-carbon double bonds, wherein one or more of the hydrogen atoms have been replaced with one or more R¹⁰⁰ groups, wherein each R¹⁰⁰ is independently a hydroxy, an oxo, a carboxyl, a carboxamido, a halo, a cyano or an amino group, as defined herein.

"Alkynyl" refers to an unsaturated acyclic C₂-C₁₀ hydrocarbon (preferably a C₂-C₈ hydrocarbon, more preferably a C₂-C₆ hydrocarbon) that can comprise one or more carbon-carbon triple bonds. Exemplary alkynyl groups include ethynyl, propynyl, butyn-1-yl, butyn-2-yl, pentyl-1-yl, pentyl-2-yl, 3-methylbutyn-1-yl, hexyl-1-yl, hexyl-2-yl, hexyl-3-yl, 3,3-dimethylbutyn-1-yl, and the like.

"Bridged cycloalkyl" refers to two or more cycloalkyl groups, heterocyclic groups, or a combination thereof fused via adjacent or non-adjacent atoms. Bridged cycloalkyl groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, hydroxy, halo, carboxyl, alkylcarboxylic acid, aryl, amidyl, ester, alkylcarboxylic ester, carboxamido, alkylcarboxamido, oxo and nitro. Exemplary bridged cycloalkyl groups include adamantyl, decahydronaphthyl, quinuclidyl, 2,6-dioxabicyclo(3.3.0)octane, 7-oxabicyclo(2.2.1)heptyl, 8-azabicyclo(3,2,1)oct-2-enyl and the like.

"Cycloalkyl" refers to a saturated or unsaturated cyclic hydrocarbon comprising from about 3 to about 10 carbon atoms. Cycloalkyl groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylarylamino, aryl, amidyl, ester, hydroxy, halo, carboxyl, alkylcarboxylic acid, alkylcarboxylic ester, carboxamido, alkylcarboxamido, oxo, alkylsulfinyl, and nitro. Exemplary cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, cyclohepta-1,3-dienyl, and the like.

"Heterocyclic ring or group" refers to a saturated or unsaturated cyclic hydrocarbon group having about 2 to about 10 carbon atoms (preferably about 4 to about 6 carbon atoms) where 1 to about 4 carbon atoms are replaced by one or more nitrogen, oxygen and/or sulfur atoms. Sulfur maybe in the thio, sulfinyl or sulfonyl oxidation state. The heterocyclic ring or group can be

fused to an aromatic hydrocarbon group. Heterocyclic groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylthio, aryloxy, arylthio, arylalkyl, hydroxy, oxo, thial, halo, carboxyl, carboxylic ester, alkylcarboxylic acid, alkylcarboxylic ester, aryl, arylcarboxylic acid, arylcarboxylic ester, amidyl, ester, alkylcarbonyl, arylcarbonyl, alkylsulfinyl, carboxamido, alkylcarboxamido, arylcarboxamido, sulfonic acid, sulfonic ester, sulfonamido and nitro. Exemplary heterocyclic groups include pyrrolyl, furyl, thienyl, 3-pyrrolyl, 4,5,6-trihydro-2H-pyran, pyridinyl, 1,4-dihydropyridinyl, pyrazolyl, triazolyl, pyrimidinyl, pyridazinyl, oxazolyl, thiazolyl, imidazolyl, indolyl, thiophenyl, furanyl, tetrahydrofuranyl, tetrazolyl, pyrrolinyl, pyrrolindinyl, oxazolindinyl, 1,3-dioxolanyl, imidazolyl, imidazolindinyl, pyrazolyl, pyrazolidinyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, 2H-pyran, 4H-pyran, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, benzo(b)thiophenyl, benzimidazolyl, benzothiazolyl, quinolinyl, and the like.

"Heterocyclic compounds" refer to mono- and polycyclic compounds comprising at least one aryl or heterocyclic ring.

"Aryl" refers to a monocyclic, bicyclic, carbocyclic or heterocyclic ring system comprising one or two aromatic rings. Exemplary aryl groups include phenyl, pyridyl, naphthyl, quinoyl, tetrahydronaphthyl, furanyl, indanyl, indenyl, indoyl, and the like. Aryl groups (including bicyclic aryl groups) can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, alkylthio, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylarylamino, halo, cyano, alkylsulfinyl, hydroxy, carboxyl, carboxylic ester, alkylcarboxylic acid, alkylcarboxylic ester, aryl, arylcarboxylic acid, arylcarboxylic ester, alkylcarbonyl, arylcarbonyl, amidyl, ester, carboxamido, alkylcarboxamido, carbomyl, sulfonic acid, sulfonic ester, sulfonamido and nitro. Exemplary substituted aryl groups include tetrafluorophenyl, pentafluorophenyl, sulfonamide, alkylsulfonyl, arylsulfonyl, and the like.

"Cycloalkenyl" refers to an unsaturated cyclic C₂-C₁₀ hydrocarbon (preferably a C₂-C₈ hydrocarbon, more preferably a C₂-C₆ hydrocarbon) which can comprise one or more carbon-carbon triple bonds.

"Alkylaryl" refers to an alkyl group, as defined herein, to which is appended an aryl group, as defined herein. Exemplary alkylaryl groups include benzyl, phenylethyl, hydroxybenzyl, fluorobenzyl, fluorophenylethyl, and the like.

"Arylalkyl" refers to an aryl radical, as defined herein, attached to an alkyl radical, as defined herein. Exemplary arylalkyl groups include benzyl, phenylethyl, 4-hydroxybenzyl, 3-fluorobenzyl, 2-fluorophenylethyl, and the like.

"Arylalkenyl" refers to an aryl radical, as defined herein, attached to an alkenyl radical, as defined herein. Exemplary arylalkenyl groups include styryl, propenylphenyl, and the like.

"Cycloalkylalkyl" refers to a cycloalkyl radical, as defined herein, attached to an alkyl radical, as defined herein.

"Cycloalkylalkoxy" refers to a cycloalkyl radical, as defined herein, attached to an alkoxy radical, as defined herein.

"Cycloalkylalkylthio" refers to a cycloalkyl radical, as defined herein, attached to an alkylthio radical, as defined herein.

"Heterocyclicalkyl" refers to a heterocyclic ring radical, as defined herein, attached to an alkyl radical, as defined herein.

"Arylheterocyclic ring" refers to a bi- or tricyclic ring comprised of an aryl ring, as defined herein, appended via two adjacent carbon atoms of the aryl ring to a heterocyclic ring, as defined herein. Exemplary arylheterocyclic rings include dihydroindole, 1,2,3,4-tetrahydroquinoline, and the like.

"Alkylheterocyclic ring" refers to a heterocyclic ring radical, as defined herein, attached to an alkyl radical, as defined herein. Exemplary alkylheterocyclic rings include 2-pyridylmethyl, 1-methylpiperidin-2-one-3-methyl, and the like.

"Alkoxy" refers to $R_{50}O-$, wherein R_{50} is an alkyl group, as defined herein (preferably a lower alkyl group or a haloalkyl group, as defined herein). Exemplary alkoxy groups include methoxy, ethoxy, t-butoxy, cyclopentyloxy, trifluoromethoxy, and the like.

"Aryloxy" refers to $R_{55}O-$, wherein R_{55} is an aryl group, as defined herein. Exemplary arylkoxy groups include naphthyloxy, quinolyloxy, isoquinolizinyloxy, and the like.

"Alkylthio" refers to $R_{50}S-$, wherein R_{50} is an alkyl group, as defined herein.

"Lower alkylthio" refers to a lower alkyl group, as defined herein, appended to a thio group, as defined herein.

"Arylalkoxy" or "alkoxyaryl" refers to an alkoxy group, as defined herein, to which is appended an aryl group, as defined herein. Exemplary arylalkoxy groups include benzyloxy, phenylethoxy, chlorophenylethoxy, and the like.

"Alkoxyalkyl" refers to an alkoxy group, as defined herein, appended to an alkyl group, as defined herein. Exemplary alkoxyalkyl groups include methoxymethyl, methoxyethyl, isopropoxymethyl, and the like.

"Alkoxyhaloalkyl" refers to an alkoxy group, as defined herein, appended to a haloalkyl group, as defined herein. Exemplary alkoxyhaloalkyl groups include 4-methoxy-2-chlorobutyl and the like.

"Cycloalkoxy" refers to $R_{54}O-$, wherein R_{54} is a cycloalkyl group or a bridged cycloalkyl group, as defined herein. Exemplary cycloalkoxy groups include cyclopropyloxy, cyclopentyloxy, cyclohexyloxy, and the like.

"Cycloalkylthio" refers to $R_{54}S-$, wherein R_{54} is a cycloalkyl group or a bridged cycloalkyl group, as defined herein. Exemplary cycloalkylthio groups include cyclopropylthio, cyclopentylthio, cyclohexylthio, and the like.

"Haloalkoxy" refers to an alkoxy group, as defined herein, in which one or more of the hydrogen atoms on the alkoxy group are substituted with halogens, as defined herein. Exemplary haloalkoxy groups include 1,1,1-trichloroethoxy, 2-bromobutoxy, and the like.

"Hydroxy" refers to $-OH$.

"Oxo" refers to $=O$.

"Oxy" refers to $-O^- R_{77}^+$ wherein R_{77} is an organic or inorganic cation.

"Oxime" refers to $=N-OR_{81}$ wherein R_{81} is a hydrogen, an alkyl group, an aryl group, an alkylsulfonyl group, an arylsulfonyl group, a carboxylic ester, an alkylcarbonyl group, an arylcarbonyl group, a carboxamido group, an alkoxyalkyl group or an alkoxyaryl group.

"Hydrazone" refers to $=N-N(R_{81})(R'_{81})$ wherein R'_{81} is independently selected from R_{81} , and R_{81} is as defined herein.

"Organic cation" refers to a positively charged organic ion. Exemplary organic cations include alkyl substituted ammonium cations, and the like.

"Inorganic cation" refers to a positively charged metal ion. Exemplary inorganic cations include Group I metal cations such as for example, sodium, potassium, magnesium, calcium, and the like.

"Hydroxyalkyl" refers to a hydroxy group, as defined herein, appended to an alkyl group, as defined herein.

"Nitrate" refers to $-O-NO_2$.

"Nitrite" refers to $-O-NO$.

5 "Thionitrate" refers to $-S-NO_2$.

"Thionitrite" and "nitrosothiol" refer to $-S-NO$.

"Nitro" refers to the group $-NO_2$ and "nitrosated" refers to compounds that have been substituted therewith.

10 "Nitroso" refers to the group $-NO$ and "nitrosylated" refers to compounds that have been substituted therewith.

"Nitrile" and "cyano" refer to $-CN$.

"Halogen" or "halo" refers to iodine (I), bromine (Br), chlorine (Cl), and/or fluorine (F).

"Amino" refers to $-NH_2$, an alkylamino group, a dialkylamino group, an arylamino group, a diarylamino group, an alkylarylamino group or a heterocyclic ring, as defined herein.

15 "Alkylamino" refers to $R_{50}NH-$, wherein R_{50} is an alkyl group, as defined herein.

Exemplary alkylamino groups include methylamino, ethylamino, butylamino, cyclohexylamino, and the like.

"Arylamino" refers to $R_{55}NH-$, wherein R_{55} is an aryl group, as defined herein.

20 "Dialkylamino" refers to $R_{52}R_{53}N-$, wherein R_{52} and R_{53} are each independently an alkyl group, as defined herein. Exemplary dialkylamino groups include dimethylamino, diethylamino, methyl propargylamino, and the like.

"Diarylamino" refers to $R_{55}R_{60}N-$, wherein R_{55} and R_{60} are each independently an aryl group, as defined herein.

25 "Alkylarylamino or arylalkylamino" refers to $R_{52}R_{55}N-$, wherein R_{52} is an alkyl group, as defined herein, and R_{55} is an aryl group, as defined herein.

"Alkylarylalkylamino" refers to $R_{52}R_{79}N-$, wherein R_{52} is an alkyl group, as defined herein, and R_{79} is an arylalkyl group, as defined herein.

"Alkylcycloalkylamino" refers to $R_{52}R_{80}N-$, wherein R_{52} is an alkyl group, as defined herein, and R_{80} is a cycloalkyl group, as defined herein.

30 "Aminoalkyl" refers to an amino group, an alkylamino group, a dialkylamino group, an arylamino group, a diarylamino group, an alkylarylamino group or a heterocyclic ring, as defined

herein, to which is appended an alkyl group, as defined herein. Exemplary aminoalkyl groups include dimethylaminopropyl, diphenylaminocyclopentyl, methylaminomethyl, and the like.

"Aminoaryl" refers to an aryl group to which is appended an alkylamino group, an arylamino group or an arylalkylamino group. Exemplary aminoaryl groups include anilino, N-methylanilino, N-benzylanilino, and the like.

"Thio" refers to $-S-$.

"Sulfinyl" refers to $-S(O)-$.

"Methanthial" refers to $-C(S)-$.

"Thial" refers to $=S$.

"Sulfonyl" refers to $-S(O)_2-$.

"Sulfonic acid" refers to $-S(O)_2OR_{76}$, wherein R_{76} is a hydrogen, an organic cation or an inorganic cation, as defined herein.

"Alkylsulfonic acid" refers to a sulfonic acid group, as defined herein, appended to an alkyl group, as defined herein.

"Arylsulfonic acid" refers to a sulfonic acid group, as defined herein, appended to an aryl group, as defined herein.

"Sulfonic ester" refers to $-S(O)_2OR_{58}$, wherein R_{58} is an alkyl group, an aryl group, or an aryl heterocyclic ring, as defined herein.

"Sulfonamido" refers to $-S(O)_2N(R_{51})(R_{57})$, wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group or an arylheterocyclic ring, as defined herein, or R_{51} and R_{57} when taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined herein.

"Alkylsulfonamido" refers to a sulfonamido group, as defined herein, appended to an alkyl group, as defined herein.

"Arylsulfonamido" refers to a sulfonamido group, as defined herein, appended to an aryl group, as defined herein.

"Alkylthio" refers to $R_{50}S-$, wherein R_{50} is an alkyl group, as defined herein (preferably a lower alkyl group, as defined herein).

"Arylthio" refers to $R_{55}S-$, wherein R_{55} is an aryl group, as defined herein.

"Arylalkylthio" refers to an aryl group, as defined herein, appended to an alkylthio group, as defined herein.

"Alkylsulfinyl" refers to $R_{50}\text{-S(O)-}$, wherein R_{50} is an alkyl group, as defined herein.

"Alkylsulfonyl" refers to $R_{50}\text{-S(O)}_2\text{-}$, wherein R_{50} is an alkyl group, as defined herein.

"Alkylsulfonyloxy" refers to $R_{50}\text{-S(O)}_2\text{-O-}$, wherein R_{50} is an alkyl group, as defined herein.

5 "Arylsulfinyl" refers to $R_{55}\text{-S(O)-}$, wherein R_{55} is an aryl group, as defined herein.

"Arylsulfonyl" refers to $R_{55}\text{-S(O)}_2\text{-}$, wherein R_{55} is an aryl group, as defined herein.

"Arylsulfonyloxy" refers to $R_{55}\text{-S(O)}_2\text{-O-}$, wherein R_{55} is an aryl group, as defined herein.

"Amidyl" refers to $R_{51}\text{C(O)N(R}_{57}\text{)-}$ wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group or an arylheterocyclic ring, as defined herein.

10 "Ester" refers to $R_{51}\text{C(O)O-}$ wherein R_{51} is a hydrogen atom, an alkyl group, an aryl group or an arylheterocyclic ring, as defined herein.

"Carbamoyl" refers to $\text{-O-C(O)N(R}_{51}\text{)(R}_{57}\text{)}$, wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group or an arylheterocyclic ring, as defined herein, or R_{51} and R_{57} taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined herein.

15 "Carboxyl" refers to -C(O)OR_{76} , wherein R_{76} is a hydrogen, an organic cation or an inorganic cation, as defined herein.

"Carbonyl" refers to -C(O)- .

"Alkylcarbonyl" refers to $R_{52}\text{-C(O)-}$, wherein R_{52} is an alkyl group, as defined herein.

20 "Arylcarbonyl" refers to $R_{55}\text{-C(O)-}$, wherein R_{55} is an aryl group, as defined herein.

"Arylalkylcarbonyl" refers to $R_{55}\text{-R}_{52}\text{-C(O)-}$, wherein R_{55} is an aryl group, as defined herein, and R_{52} is an alkyl group, as defined herein.

"Alkylarylcarbonyl" refers to $R_{52}\text{-R}_{55}\text{-C(O)-}$, wherein R_{55} is an aryl group, as defined herein, and R_{52} is an alkyl group, as defined herein.

25 "Heterocyclicalkylcarbonyl" refer to $R_{78}\text{C(O)-}$ wherein R_{78} is a heterocyclicalkyl group, as defined herein.

"Carboxylic ester" refers to -C(O)OR_{58} , wherein R_{58} is an alkyl group, an aryl group or an aryl heterocyclic ring, as defined herein.

30 "Alkylcarboxylic acid" and "alkylcarboxyl" refer to an alkyl group, as defined herein, appended to a carboxyl group, as defined herein.

"Alkylcarboxylic ester" refers to an alkyl group, as defined herein, appended to a

carboxylic ester group, as defined herein.

"Arylcarboxylic acid" refers to an aryl group, as defined herein, appended to a carboxyl group, as defined herein.

5 "Arylcarboxylic ester" and "arylcarboxyl" refer to an aryl group, as defined herein, appended to a carboxylic ester group, as defined herein.

"Carboxamido" refers to $-C(O)N(R_{51})(R_{57})$, wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group or an arylheterocyclic ring, as defined herein, or R_{51} and R_{57} when taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined herein.

10 "Alkylcarboxamido" refers to an alkyl group, as defined herein, appended to a carboxamido group, as defined herein.

"Arylcarboxamido" refers to an aryl group, as defined herein, appended to a carboxamido group, as defined herein.

15 "Urea" refers to $-N(R_{59})-C(O)N(R_{51})(R_{57})$ wherein R_{51} , R_{57} , and R_{59} are each independently a hydrogen atom, an alkyl group, an aryl group or an arylheterocyclic ring, as defined herein, or R_{51} and R_{57} taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined herein.

20 "Phosphoryl" refers to $-P(R_{70})(R_{71})(R_{72})$, wherein R_{70} is a lone pair of electrons, thial or oxo, and R_{71} and R_{72} are each independently a covalent bond, a hydrogen, a lower alkyl, an alkoxy, an alkylamino, a hydroxy, an oxy or an aryl, as defined herein.

"Silyl" refers to $-Si(R_{73})(R_{74})(R_{75})$, wherein R_{73} , R_{74} and R_{75} are each independently a covalent bond, a lower alkyl, an alkoxy, an aryl or an arylalkoxy, as defined herein.


25 The cardiovascular compounds used in the compounds and compositions of the invention are preferably aldosterone antagonists, angiotensin II antagonists, hydralazine compounds, calcium channel blockers, diuretics, endothelin antagonists, neutral endopeptidase inhibitors and renin inhibitors.

Suitable aldosterone antagonists include, but are not limited to, canrenone, potassium canrenoate, spironolactone, eplerenone, and the like.

30 Suitable angiotensin II antagonists, include, but are not limited to, angiotensin, candesartan, cilexetil, eprosartan, irbesartan, losartan, olmesartan, medoxomil, remikirin saralasin, tasosartan, telmisartan, valsartan, zolasartin, BMS 184698, 3-(2'(tetrazole-5-yl)-1,1'-

biphen-4-yl)methyl-5,7-dimethyl-2-ethyl-3H-imidazo(4,5-b)pyridine, antibodies to angiotensin II, and the like.

Suitable hydralazine compounds include, but are not limited to, compounds having the formula:



wherein a, b and c are independently a single or double bond; R₁ and R₂ are each independently a hydrogen, an alkyl, an ester or a heterocyclic ring, wherein alkyl, ester and heterocyclic ring are as defined herein; R₃ and R₄ are each independently a lone pair of electrons or a hydrogen, with the proviso that at least one of R₁, R₂, R₃ and R₄ is not a hydrogen.

10 Exemplary hydralazine compounds include budralazine, cadralazine, dihydralazine, endralazine, hydralazine, pildralazine, todralazine, and the like.

Suitable calcium channel blockers, include, but are not limited to, amlodipine, aranidipine, barnidipine, benidipine, bepridil, cilnidipine, cinnarizine, clentiazem, diltiazem, dotarizine, efonidipine, elgodipine, fantofarone, felodipine, flunarizine, fluspirilene, gallopamil, isradipine, lacidipine, lercanidipine, lomerizine, manidipine, mibefradil, monatepil, nicardipine, nifedipine, nilvadipine, nimodipine, nisoldipine, nitrendipine, semotiadil, verapamil, and the like. Suitable calcium channel blockers are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

20 Suitable diuretics include, but are not limited to, thiazides (such as, for example, althiazide, bendroflumethiazide, benzclortriazide, benzthiazide, buthiazide, chlorothiazide, cyclophenethiazide, cyclothiazide, ethiazide, hydrochlorothiazide, methyclothiazide, penflutazide, polythiazide, teclothiazide, trichlormethiazide, triflumethazide, and the like); ambuside, amiloride, aminometradine, azosemide, bemetizide, bumetanide, butazolamide, butizide, ethacrynic acid, canrenone, chloraminophenamide, chlorazanil, chlormerodrin, chlorthalidone, clofenamide, clopamide, clorexolone, disulfamide, ethacrynic acid, ethoxzolamide, etozolon, fenquizone, furosemide, mefruside, meralluride, mercaptomerin sodium, mercumallylic acid, mersalyl, methazolamide, metolazone, muzolimine, pamabrom,

paraflutizide, piretanide, protheobromine, quinethazone, scoparius, spironalactone, theobromine, ticrynafen, torsemide, triamterene, xipamide or potassium, and the like.

Suitable endothelin antagonists, include, but are not limited to, bosentan, endothelin, sulfonamide endothelin antagonists, BQ-123, SQ 28608, and the like.

5 Suitable neutral endopeptidase inhibitors, include, but are not limited to, atrial natriuretic peptides, diazapins, azepinones, ecadotril, omapatrilat, sampatrilat, BMS 189,921, and the like.

 Suitable renin inhibitors, include, but are not limited to, aldosterone, enalkrein, medullipin, tonin, RO 42-5892, A 65317, CP 80794, ES 1005, ES 8891, SQ 34017, urea derivatives of peptides, amino acids connected by nonpeptide bonds, di- and tri-peptide
10 derivatives, amino acids and derivatives thereof, diol sulfonamides and sulfinyls, modified peptides, peptidyl beta-aminoacyl aminodiols, carbamates, monoclonal antibodies to renin, and the like. Suitable renin inhibitors are described more fully in U.S. Patent Nos. 5,116,835, 5,114,937, 5,106,835, 5,104,869, 5,095,119, 5,098,924, 5,095,006, 5,089,471, 5,075,451, 5,066,643, 5,063,208, 4,845,079, 5,055,466, 4,980,283, 4,885,292, 4,780,401, 5,071,837, 5,064,965,
15 5,063,207, 5,036,054, 5,036,053, 5,034,512, and 4,894,437, the disclosures of each of which are incorporated herein by reference in their entirety;

 The contemplated cardiovascular compounds of the invention are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, (1996); Merck Index on CD-ROM, 13th Edition; STN Express, file phar
20 and file registry, the disclosures of each of which are incorporated by reference herein in their entirety.

 In one embodiment the cardiovascular compounds of the invention are aldosterone antagonists, angiotensin II antagonists, hydralazine compounds, calcium channel blockers, diuretics, endothelin antagonists, neutral endopeptidase inhibitors and renin inhibitors, that
25 contain one or more of the following functionalities: a carboxylic acid group (-COOH), a hydroxyl group (-OH), a thiol group (-SH) and/or a primary or secondary amine group (-NH). The cardiovascular compounds of the invention are nitrosated and/or nitrosylated through one or more of these functionalities such as oxygen (hydroxyl condensation), sulfur (sulfhydryl condensation) and/or nitrogen.

30 In one embodiment, the invention describes nitrosated and/or nitrosylated cardiovascular compounds of the invention that are nitrosated and/or nitrosylated aldosterone antagonists,

nitrosated and/or nitrosylated angiotensin II antagonists, nitrosated and/or nitrosylated hydralazine compounds, nitrosated and/or nitrosylated calcium channel blockers, nitrosated and/or nitrosylated diuretics, nitrosated and/or nitrosylated endothelin antagonists, nitrosated and/or nitrosylated neutral endopeptidase inhibitors and nitrosated and/or nitrosylated renin inhibitors where the cardiovascular compounds of the invention must contain at least one nitrosated and/or nitrosylated carboxylic acid group, nitrosated and/or nitrosylated hydroxyl group, nitrosated and/or nitrosylated thiol group and/or primary or secondary nitrosated and/or nitrosylated amine group. In another embodiment, the cardiovascular compounds are nitrosated and/or nitrosylated by containing the group K;

wherein

K is $-W_a-E_b-(C(R_e)(R_f))_p-E_c-(C(R_e)(R_f))_x-W_d-(C(R_e)(R_f))_y-W_i-E_j-W_g-(C(R_e)(R_f))_z-T-Q$;

a, b, c, d, g, i and j are each independently an integer from 0 to 3;

p, x, y and z are each independently an integer from 0 to 10;

Q is $-NO$ or $-NO_2$;

W at each occurrence is independently $-C(O)-$, $-C(S)-$, $-T-$, $-(C(R_e)(R_f))_h-$, an alkyl group, an aryl group, a heterocyclic ring, an arylheterocyclic ring, or $-(CH_2CH_2O)_q-$;

E at each occurrence is independently $-T-$, an alkyl group, an aryl group, $-(C(R_e)(R_f))_h-$, a heterocyclic ring, an arylheterocyclic ring, or $-(CH_2CH_2O)_q-$;

h is an integer from 1 to 10;

q is an integer from 1 to 5;

R_e and R_f are each independently a hydrogen, an alkyl, a cycloalkoxy, a halogen, a hydroxy, an hydroxyalkyl, an alkoxyalkyl, an arylheterocyclic ring, an alkylaryl, an alkylcycloalkyl, an alkylheterocyclic ring, a cycloalkylalkyl, a cycloalkylthio, a cycloalkenyl, an heterocyclicalkyl, an alkoxy, a haloalkoxy, an amino, an alkylamino, a dialkylamino, an arylamino, a diarylamino, an alkylaryl amino, an alkoxyhaloalkyl, a sulfonic acid, a sulfonic ester, an alkylsulfonic acid, an arylsulfonic acid, an arylalkoxy, an alkylthio, an arylthio, a cyano an aminoalkyl, an aminoaryl, an aryl, an arylalkyl, an alkylaryl, a carboxamido, an alkylcarboxamido, an arylcarboxamido, an amidyl, a carboxyl, a carbamoyl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarbonyl, an arylcarbonyl, an ester, a carboxylic ester, an alkylcarboxylic ester, an arylcarboxylic ester, a sulfonamido, an alkylsulfonamido, an arylsulfonamido, an alkylsulfonyl, an alkylsulfonyloxy, an arylsulfonyl, arylsulphonyloxy, a

sulfonic ester, a urea, a phosphoryl, a nitro, W_h , $-T-Q$, or $-(C(R_g)(R_h))_k-T-Q$, or R_e and R_f taken together with the carbons to which they are attached form a carbonyl, a methanthial, a heterocyclic ring, a cycloalkyl group, an aryl group, an oxime, a hydrazone or a bridged cycloalkyl group;

5 R_g and R_h at each occurrence are independently R_e ;

k is an integer from 1 to 3;

T at each occurrence is independently a covalent bond, a carbonyl, an oxygen, $-S(O)_o-$ or $-N(R_a)R_i-$;

o is an integer from 0 to 2;

10 R_a is a lone pair of electrons, a hydrogen or an alkyl group;

R_i is a hydrogen, an alkyl, an aryl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarboxylic ester, an arylcarboxylic ester, an alkylcarboxamido, an arylcarboxamido, an alkylaryl, an alkylsulfinyl, an alkylsulfonyl, an alkylsulfonyloxy, an arylsulfinyl, an arylsulfonyl, arylsulphonyloxy, a sulfonamido, a carboxamido, a carboxylic ester, an aminoalkyl, an aminoaryl, $-CH_2-C(T-Q)(R_e)(R_f)$, a bond to an adjacent atom creating a double bond to that atom, $-(N_2O_2-)^- \cdot M^+$, wherein M^+ is an organic or inorganic cation; and

15 with the proviso that the nitrosated and/or nitrosylated cardiovascular compounds of the invention must contain least one of a nitrite, a nitrate, a thionitrite group or a thionitrate group.

In cases where multiple designations of variables which reside in sequence are chosen as a "covalent bond" or the integer chosen is 0, the intent is to denote a single covalent bond connecting one radical to another. For example, E_0 would denote a covalent bond, while E_2 denotes $(E-E)$ and $(C(R_4)(R_4))_2$ denotes $-C(R_4)(R_4)-C(R_4)(R_4)-$.

20 Compounds of the invention that have one or more asymmetric carbon atoms may exist as the optically pure enantiomers, pure diastereomers, mixtures of enantiomers, mixtures of diastereomers, racemic mixtures of enantiomers, diastereomeric racemates or mixtures of diastereomeric racemates. It is to be understood that the invention anticipates and includes within its scope all such isomers and mixtures thereof.

25 Another embodiment of the invention describes the metabolites of the nitrosated and/or nitrosylated cardiovascular compounds and pharmaceutically acceptable salts thereof. These metabolites, include but are not limited to, the non-nitrosated and/or nitrosylated derivatives, degradation products, hydrolysis products, and the like, of the nitrosated and/or nitrosylated

cardiovascular compounds and pharmaceutically acceptable salts thereof.

Another embodiment of the invention provides processes for making the novel compounds of the invention and to the intermediates useful in such processes. The reactions are performed in solvents appropriate to the reagents and materials used are suitable for the transformations being effected. It is understood by one skilled in the art of organic synthesis that the functionality present in the molecule must be consistent with the chemical transformation proposed. This will, on occasion, necessitate judgment by the routineer as to the order of synthetic steps, protecting groups required, and deprotection conditions. Substituents on the starting materials may be incompatible with some of the reaction conditions required in some of the methods described, but alternative methods and substituents compatible with the reaction conditions will be readily apparent to one skilled in the art. The use of sulfur and oxygen protecting groups is well known for protecting thiol and alcohol groups against undesirable reactions during a synthetic procedure and many such protecting groups are known and described by, for example, Greene and Wuts, *Protective Groups in Organic Synthesis*, Third Edition, John Wiley & Sons, New York (1999).

The chemical reactions described herein are generally disclosed in terms of their broadest application to the preparation of the compounds of this invention. Occasionally, the reactions may not be applicable as described to each compound included within the disclosed scope. The compounds for which this occurs will be readily recognized by one skilled in the art. In all such cases, either the reactions can be successfully performed by conventional modifications known to one skilled in the art, *e.g.*, by appropriate protection of interfering groups, by changing to alternative conventional reagents, by routine modification of reaction conditions, and the like, or other reactions disclosed herein or otherwise conventional, will be applicable to the preparation of the corresponding compounds of this invention. In all preparative methods, all starting materials are known or readily prepared from known starting materials.

The cardiovascular compounds are nitrosated and/or nitrosylated through one or more sites such as oxygen, sulfur and/or nitrogen using conventional methods known to one skilled in the art. For example, known methods for nitrosating and/or nitrosylating compounds are described in U.S. Patent Nos. 5,380,758, 5,859,053, 5,703,073 and 6,297,260; and in WO 94/03421, WO 94/04484, WO 94/12463, WO 95/09831, WO 95/19952, WO 95/30641, WO 97/27749, WO 98/19672, WO 98/21193, WO 00/51988, WO 00/61604, WO 00/72838, WO

01/00563, WO 01/04082, WO 01/10814, WO 01/12584, WO 01/45703, WO 00/61541, WO 00/61537, WO 02/11707, WO 02/30866 and in Oae et al, *Org. Prep. Proc. Int.*, 15(3):165-198 (1983), the disclosures of each of which are incorporated by reference herein in their entirety.

The methods of nitrosating and/or nitrosylating the compounds described in these references can be applied by one skilled in the art to produce any of the nitrosated and/or nitrosylated cardiovascular compounds described herein. The nitrosated and/or nitrosylated cardiovascular compounds of the invention donate, transfer or release a biologically active form of nitrogen monoxide (i.e., nitric oxide).

Nitrogen monoxide can exist in three forms: NO⁻ (nitroxyl), NO[•] (uncharged nitric oxide) and NO⁺ (nitrosonium). NO[•] is a highly reactive short-lived species that is potentially toxic to cells. This is critical because the pharmacological efficacy of NO depends upon the form in which it is delivered. In contrast to the nitric oxide radical (NO[•]), nitrosonium (NO⁺) does not react with O₂ or O₂⁻ species, and functionalities capable of transferring and/or releasing NO⁺ and NO⁻ are also resistant to decomposition in the presence of many redox metals. Consequently, administration of charged NO equivalents (positive and/or negative) is a more effective means of delivering a biologically active NO to the desired site of action.

Compounds contemplated for use in the invention, e.g., cardiovascular compounds that are nitrosated and/or nitrosylated, through one or more sites such as oxygen (hydroxyl condensation), sulfur (sulfhydryl condensation) and/or nitrogen, are, optionally, used in combination with nitric oxide and compounds that release nitric oxide or otherwise directly or indirectly deliver or transfer a biologically active form of nitrogen monoxide to a site of its intended activity, such as on a cell membrane *in vivo*.

The term "nitric oxide" encompasses uncharged nitric oxide (NO[•]) and charged nitrogen monoxide species, preferably charged nitrogen monoxide species, such as nitrosonium ion (NO⁺) and nitroxyl ion (NO⁻). The reactive form of nitric oxide can be provided by gaseous nitric oxide. The nitrogen monoxide releasing, delivering or transferring compounds have the structure F-NO, wherein F is a nitrogen monoxide releasing, delivering or transferring moiety, and include any and all such compounds which provide nitrogen monoxide to its intended site of action in a form active for its intended purpose. The term "NO adducts" encompasses any nitrogen monoxide releasing, delivering or transferring compounds, including, for example, S-nitrosothiols, nitrites, nitrates, S-nitrothiols, sydnonimines, 2-hydroxy-2-nitrosohydrazines,

(NONOates), (E)-alkyl-2-((E)-hydroxyimino)-5-nitro-3-hexeneamide (FK-409), (E)-alkyl-2-((E)-hydroxyimino)-5-nitro-3-hexeneamines, N-((2Z, 3E)-4-ethyl-2-(hydroxyimino)-6-methyl-5-nitro-3-heptenyl)-3-pyridinecarboxamide (FR 146801), nitrosoamines, furoxans as well as substrates for the endogenous enzymes which synthesize nitric oxide. NONOates include, but are not limited to, (Z)-1-(N-methyl-N-(6-(N-methyl-ammoniohexyl)amino))diazene-1-ium-1,2-diolate ("MAHMA/NO"), (Z)-1-(N-(3-ammoniopropyl)-N-(n-propyl)amino)diazene-1-ium-1,2-diolate ("PAPA/NO"), (Z)-1-(N-(3-aminopropyl)-N-(4-(3-aminopropylammonio)butyl)-amino)diazene-1-ium-1,2-diolate (spermine NONOate or "SPER/NO") and sodium(Z)-1-(N,N-diethylamino)diazonium-1,2-diolate (diethylamine NONOate or "DEA/NO") and derivatives thereof. NONOates are also described in U.S. Patent Nos. 6,232,336, 5,910,316 and 5,650,447, the disclosures of which are incorporated herein by reference in their entirety. The "NO adducts" can be mono-nitrosylated, poly-nitrosylated, mono-nitrosated and/or poly-nitrosated at a variety of naturally susceptible or artificially provided binding sites for biologically active forms of nitrogen monoxide.

One group of NO adducts is the S-nitrosothiols, which are compounds that include at least one -S-NO group. These compounds include S-nitroso-polypeptides (the term "polypeptide" includes proteins and polyamino acids that do not possess an ascertained biological function, and derivatives thereof); S-nitrosylated amino acids (including natural and synthetic amino acids and their stereoisomers and racemic mixtures and derivatives thereof); S-nitrosylated sugars; S-nitrosylated, modified and unmodified, oligonucleotides (preferably of at least 5, and more preferably 5-200 nucleotides); straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted S-nitrosylated hydrocarbons; and S-nitroso heterocyclic compounds. S-nitrosothiols and methods for preparing them are described in U.S. Patent Nos. 5,380,758 and 5,703,073; WO 97/27749; WO 98/19672; and Oae et al, *Org. Prep. Proc. Int.*, 15(3):165-198 (1983), the disclosures of each of which are incorporated by reference herein in their entirety.

Another embodiment of the invention is S-nitroso amino acids where the nitroso group is linked to a sulfur group of a sulfur-containing amino acid or derivative thereof. Such compounds include, for example, S-nitroso-N-acetylcysteine, S-nitroso-captopril, S-nitroso-N-acetylpenicillamine, S-nitroso-homocysteine, S-nitroso-cysteine, S-nitroso-glutathione, S-nitroso-cysteinyl-glycine, and the like.

Suitable S-nitrosylated proteins include thiol-containing proteins (where the NO group is attached to one or more sulfur groups on an amino acid or amino acid derivative thereof) from various functional classes including enzymes, such as tissue-type plasminogen activator (TPA) and cathepsin B; transport proteins, such as lipoproteins; heme proteins, such as hemoglobin and serum albumin; and biologically protective proteins, such as immunoglobulins, antibodies and cytokines. Such nitrosylated proteins are described in WO 93/09806, the disclosure of which is incorporated by reference herein in its entirety. Examples include polynitrosylated albumin where one or more thiol or other nucleophilic centers in the protein are modified.

Other examples of suitable S-nitrosothiols include:

- (i) $\text{HS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{SNO}$;
- (ii) $\text{ONS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{R}_e$; or
- (iii) $\text{H}_2\text{N}-\text{CH}(\text{CO}_2\text{H})-(\text{CH}_2)_m-\text{C}(\text{O})\text{NH}-\text{CH}(\text{CH}_2\text{SNO})-\text{C}(\text{O})\text{NH}-\text{CH}_2-\text{CO}_2\text{H}$;

wherein m is an integer from 2 to 20; R_e and R_f are each independently a hydrogen, an alkyl, a cycloalkoxy, a halogen, a hydroxy, an hydroxyalkyl, an alkoxyalkyl, an arylheterocyclic ring, a cycloalkylalkyl, a heterocyclicalkyl, an alkoxy, a haloalkoxy, an amino, an alkylamino, a dialkylamino, an arylamino, a diarylamino, an alkylarylamino, an alkoxyhaloalkyl, a haloalkoxy, a sulfonic acid, a sulfonic ester, an alkylsulfonic acid, an arylsulfonic acid, an arylalkoxy, an alkylthio, an arylthio, a cyano, an aminoalkyl, an aminoaryl, an aryl, an arylalkyl, a carboxamido, an alkylcarboxamido, an arylcarboxamido, an amidyl, a carboxyl, a carbamoyl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarbonyl, an arylcarbonyl, an ester, a carboxylic ester, an alkylcarboxylic ester, an arylcarboxylic ester, a haloalkoxy, a sulfonamido, an alkylsulfonamido, an arylsulfonamido, an alkylsulfonyl, an alkylsulfonyloxy, an arylsulfonyl, an arylsulfonyloxy, a urea, a nitro, -T-Q-, or $-(\text{C}(\text{R}_g)(\text{R}_h))_k-\text{T}-\text{Q}$ or R_e and R_f taken together are an oxo, a methanthial, a heterocyclic ring, a cycloalkyl group, an oxime, a hydrazone or a bridged cycloalkyl group; Q is -NO or -NO₂; and T is independently a covalent bond, a carbonyl, an oxygen, -S(O)_o- or -N(R_a)R_i, wherein o is an integer from 0 to 2, R_a is a lone pair of electrons, a hydrogen or an alkyl group; R_i is a hydrogen, an alkyl, an aryl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarboxylic ester, an arylcarboxylic ester, an alkylcarboxamido, an arylcarboxamido, an alkylsulfinyl, an alkylsulfonyl, an alkylsulfonyloxy, an arylsulfinyl, an arylsulfonyloxy, an arylsulfonyl, a sulfonamido, a carboxamido, a carboxylic ester, an aminoalkyl, an aminoaryl, -CH₂-C(T-Q)(R_g)(R_h), or $-(\text{N}_2\text{O}_2)^-\cdot\text{M}^+$, wherein M⁺ is an organic or inorganic cation; with the

proviso that when R_i is $-\text{CH}_2-\text{C}(\text{T}-\text{Q})(\text{R}_g)(\text{R}_h)$ or $-(\text{N}_2\text{O}_2-)\cdot\text{M}^+$; then "-T-Q" can be a hydrogen, an alkyl group, an alkoxyalkyl group, an aminoalkyl group, a hydroxy group or an aryl group; and

R_g and R_h at each occurrence are independently R_e ;

In cases where R_e and R_f are a heterocyclic ring or taken together R_e and R_f are a heterocyclic ring, then R_i can be a substituent on any disubstituted nitrogen contained within the radical wherein R_i is as defined herein.

Nitrosothiols can be prepared by various methods of synthesis. In general, the thiol precursor is prepared first, then converted to the S-nitrosothiol derivative by nitrosation of the thiol group with NaNO_2 under acidic conditions (pH is about 2.5) which yields the S-nitroso derivative. Acids which can be used for this purpose include aqueous sulfuric, acetic and hydrochloric acids. The thiol precursor can also be nitrosylated by reaction with an organic nitrite such as tert-butyl nitrite, or a nitrosonium salt such as nitrosonium tetrafluoroborate in an inert solvent.

Another group of NO adducts for use in the invention, where the NO adduct is a compound that donates, transfers or releases nitric oxide, include compounds comprising at least one ON-O- or ON-N- group. The compounds that include at least one ON-O- or ON-N- group are preferably ON-O- or ON-N-polypeptides (the term "polypeptide" includes proteins and polyamino acids that do not possess an ascertained biological function, and derivatives thereof); ON-O- or ON-N-amino acids (including natural and synthetic amino acids and their stereoisomers and racemic mixtures); ON-O- or ON-N-sugars; ON-O- or -ON-N- modified or unmodified oligonucleotides (comprising at least 5 nucleotides, preferably 5-200 nucleotides); ON-O- or ON-N- straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbons; and ON-O-, ON-N- or ON-C-heterocyclic compounds.

Another group of NO adducts for use in the invention include nitrates that donate, transfer or release nitric oxide, such as compounds comprising at least one $\text{O}_2\text{N-O-}$, $\text{O}_2\text{N-N-}$ or $\text{O}_2\text{N-S-}$ group. Preferred among these compounds are $\text{O}_2\text{N-O-}$, $\text{O}_2\text{N-N-}$ or $\text{O}_2\text{N-S-}$ polypeptides (the term "polypeptide" includes proteins and also polyamino acids that do not possess an ascertained biological function, and derivatives thereof); $\text{O}_2\text{N-O-}$, $\text{O}_2\text{N-N-}$ or $\text{O}_2\text{N-S-}$ amino acids (including natural and synthetic amino acids and their stereoisomers and racemic mixtures); $\text{O}_2\text{N-O-}$, $\text{O}_2\text{N-N-}$ or $\text{O}_2\text{N-S-}$ sugars; $\text{O}_2\text{N-O-}$, $\text{O}_2\text{N-N-}$ or $\text{O}_2\text{N-S-}$ modified and unmodified oligonucleotides (comprising at least 5 nucleotides, preferably 5-200 nucleotides);

O₂N-O-, O₂N-N- or O₂N-S- straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbons; and O₂N-O-, O₂N-N- or O₂N-S- heterocyclic compounds. Preferred examples of compounds comprising at least one O₂N-O-, O₂N-N- or O₂N-S- group include isosorbide dinitrate, isosorbide mononitrate, clonitrate, erythrityl tetranitrate, mannitol hexanitrate, nitroglycerin, pentaerythritoltetranitrate, pentrinitrol, propatynitrate and organic nitrates with a sulfhydryl-containing amino acid such as, for example SPM 3672, SPM 5185, SPM 5186 and those disclosed in U. S. Patent Nos. 5,284,872, 5,428,061, 5,661,129, 5,807,847 and 5,883,122 and in WO 97/46521, WO 00/54756 and in WO 03/013432, the disclosures of each of which are incorporated by reference herein in their entirety.

Another group of NO adducts are N-oxo-N-nitrosoamines that donate, transfer or release nitric oxide and are represented by the formula: R¹R²N-N(O-M⁺)-NO, where R¹ and R² are each independently a polypeptide, an amino acid, a sugar, a modified or unmodified oligonucleotide, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbon, or a heterocyclic group, and where M⁺ is an organic or inorganic cation, such as, for example, an alkyl substituted ammonium cation or a Group I metal cation.

The invention is also directed to compounds that stimulate endogenous NO or elevate levels of endogenous endothelium-derived relaxing factor (EDRF) *in vivo* or are substrates for nitric oxide synthase. Such compounds include, for example, L-arginine, L-homoarginine, and N-hydroxy-L-arginine, including their nitrosated and/or nitrosylated analogs (e.g., nitrosated L-arginine, nitrosylated L-arginine, nitrosated N-hydroxy-L-arginine, nitrosylated N-hydroxy-L-arginine, nitrosated L-homoarginine and nitrosylated L-homoarginine), precursors of L-arginine and/or physiologically acceptable salts thereof, including, for example, citrulline, ornithine, glutamine, lysine, polypeptides comprising at least one of these amino acids, inhibitors of the enzyme arginase (e.g., N-hydroxy-L-arginine and 2(S)-amino-6-boronoheptanoic acid), nitric oxide mediators and/or physiologically acceptable salts thereof, including, for example, pyruvate, pyruvate precursors, α -keto acids having four or more carbon atoms, precursors of α -keto acids having four or more carbon atoms (as disclosed in WO 03/017996, the disclosure of which is incorporated herein in its entirety), and the substrates for nitric oxide synthase, cytokines, adenosin, bradykinin, calreticulin, bisacodyl, and phenolphthalein. EDRF is a vascular relaxing factor secreted by the endothelium, and has been identified as nitric oxide (NO) or a closely related derivative thereof (Palmer et al, *Nature*, 327:524-526 (1987); Ignarro et al, *Proc. Natl.*

Acad. Sci. USA, 84:9265-9269 (1987)).

The invention is also based on the discovery that compounds and compositions of the invention may be used in conjunction with other therapeutic agents for co-therapies, partially or completely, in place of other therapeutic agents, such as, for example, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H₂ receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and mixtures of two or more thereof. The therapeutic agent may optionally be nitrosated and/or nitrosylated.

Suitable aldosterone antagonists include, but are not limited to, canrenone, potassium canrenoate, spironolactone, eplerenone, and the like. Suitable aldosterone antagonists are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable alpha-adrenergic receptor antagonists include but are not limited to, phentolamine, tolazoline, idazoxan, deriglidole, RX 821002, BRL 44408, BRL 44409, BAM 1303, labetalol, ifenprodil, rauwolscine, corynathine, raubasine, tetrahydroalstonine, apoyohimbine, akuammigine, β -yohimbine, yohimbol, yohimbine, pseudoyohimbine, epi-3 α -yohimbine, 10-hydroxy-yohimbine, 11-hydroxy-yohimbine, tamsulosin, benoxathian, atipamezole, BE 2254, WB 4101, HU-723, tedisamil, mirtazipine, setiptiline, reboxitine, delequamine, naftopil, saterinone, SL 89.0591, ARC 239, urapidil, 5-methylurapidil, monatepi, haloperidol, indoramin, SB 216469, moxislyte, trazodone, dapiprozole, efaroxan, Recordati 15/2739, SNAP 1069, SNAP 5089, SNAP 5272, RS 17053, SL 89.0591, KMD 3213, spiperone, AH 11110A, chloroethylclonidine, BMY 7378, niguldipine, and the like. Suitable alpha-adrenergic receptor antagonists are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable antidiabetic compounds include but are not limited to, acarbose, acetoexamide, buformin, carbutamide, chlorpropamide, glibornuride, gliclazide, glimepiride, glipizide, gliquidone, glisoxepid, glyburide, glybutiazol(e), glybuzole, glyhexamide, glymidine, glypinamide, insulin, metformin, miglitol, nateglinide, phenbutamide, phenformin, pioglitazone, repaglinide, rosiglitazone, tolazamide, tolbutamide, tolcyclamide, troglitazone, voglibose, and the like. Suitable antidiabetic compounds are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable anti-hyperlipidemic drugs include, but are not limited to, statins or HMG-CoA reductase inhibitors, such as, for example, atorvastatin (LIPITOR®), cerivastatin (BAYCOL®), fluvastatin (Sandoz XU-62-320), fluvastatin, lovastatin (MEVACOR®), mevastatin, pravastatin (PRAVACHOL®), simvastatin (ZOCOR®), velostatin (also known as synvinolin) and the like; gemfibrozil, cholestyramine, colestipol, nicotinic acid, bile acid sequestrants, such as, for example, cholestyramine, colessevelam, colestipol, poly(methyl-(3-trimethylaminopropyl) imino-trimethylene dihalide) and the like; probucol; fibric acid agents or fibrates, such as, for example, bezafibrate (Bezalip™), beclobrate, binifibrate, ciprofibrate, clinofibrate, clofibrate, etofibrate, fenofibrate (Lipidil™, Lipidil Micro™), gemfibrozil (Lopid™.), nicofibrate, pirifibrate, ronifibrate, simfibrate, theofibrate and the like.

Suitable angiotensin II antagonists include, but are not limited to, angiotensin, candesartan, cilexetil, eprosartan, irbesartan, losartan, olmesartan, medoxomil, remikirin saralasin, tasosartan, telmisartan, valsartan, zolasartin, BMS 184698, 3-(2'-(tetrazole-5-yl)-1,1'-biphen-4-yl)methyl-5,7-dimethyl-2-ethyl-3H-imidazo(4,5-b)pyridine, antibodies to angiotensin II, and the like. Suitable angiotensin II antagonists are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable angiotensin-converting enzyme inhibitors (ACE inhibitors) include, but are not limited to, alacepril, benazepril, benazeprilat, captopril, ceronapril, cilazapril, delapril, duinapril, enalapril, enalaprilat, fosinopril, imidapril, lisinopril, losartan, moveltipril, moexipril, naphthopidil, pentopril, perindopril, quinapril, ramipril, rentipril, spirapril, temocapril,

trandolapril, urapidil, zofenopril, acylmercapto and mercaptoalkanoyl pralines, carboxyalkyl dipeptides, carboxyalkyl dipeptide, phosphinylalkanoyl pralines, and the like.

Suitable antioxidants include, but are not limited to, small-molecule antioxidants and antioxidant enzymes. Suitable small-molecule antioxidants include, but are not limited to, 5 hydralazine compounds, glutathione, vitamin C, vitamin E, cysteine, N-acetyl-cysteine, β -carotene, ubiquinone, ubiquinol-10, tocopherols, coenzyme Q, superoxide dismutase mimetics and the like. Suitable antioxidant enzymes include, but are not limited to, superoxide dismutase, catalase, glutathione peroxidase, and the like. The antioxidant enzymes can be delivered by gene therapy as a viral vector and/or a non-viral vector. Suitable antioxidants are 10 described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable antithrombotic and vasodilator drugs include, but are not limited to, acetorphan, acetylsalicylic acid, argatroban, bamethan, benfurodil, benziodarone, betahistine, 15 brovincamine, bufenide, citicoline, clobenfurol, clopidogrel, cyclandelate, dalteparin, dipyridamol, droprenilamine, enoxaparin, fendiline, ifenprodil, iloprost, indobufen, isobogrel, isoxsuprine, heparin, lamifiban, midrodine, nadroparin, nicotinoyl alcohol, nylidrin, ozagrel, perhexiline, phenylpropanolamine, prenylamine, papaveroline, reviparin sodium salt, ridogrel, suloctidil, tinofedrine, tinzaparin, trifusal, xanthinal niacinate, and the like. Suitable 20 antithrombotic and vasodilator drugs are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable beta-adrenergic blockers include, but are not limited to, acebutolol, alprenolol, 25 amosulalol, arotinolol, atenolol, befunolol, betaxolol, bevantolol, bisoprolol, bopindolol, bucindolol, bucumolol, bufetolol, bufuralol, bunitrolol, bupranolol, butafilolol, carazolol, carteolol, carvedilol, celiprolol, cetamolol, cindolol, cloranolol, dilevalol, epanolol, esmolol, indenolol, labetalol, landiolol, mepindolol, metipranolol, metoprolol, moprolol, nadolol, nadoxolol, nebivolol, nifenalol, nipradilol, oxprenolol, penbutolol, pindolol, practolol, 30 pronethalol, propranolol, sotalol, sulfinalol, talinolol, tertatolol, tilisolol, timolol, toliprolol, xibenolol, and the like. Suitable beta-adrenergic blockers are described more fully in the

literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable calcium channel blockers include, but are not limited to, amlodipine, aranidipine, barnidipine, benidipine, bepridil, cilnidipine, cinnarizine, clentiazem, diltiazem, dotarizine, efonidipine, elgodipine, fantofarone, felodipine, flunarizine, fluspirilene, gallopamil, isradipine, lacidipine, lercanidipine, lomerizine, manidipine, mibefradil, monatepil, nifedipine, nifedipine, nilvadipine, nimodipine, nisoldipine, nitrendipine, semotiadil, verapamil, and the like. Suitable calcium channel blockers are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable diuretics include but are not limited to, thiazides (such as, for example, althiazide, bendroflumethiazide, benzclortriazide, benzthiazide, buthiazide, chlorothiazide, cyclopenethiazide, cyclothiazide, ethiazide, hydrochlorothiazide, methyclothiazide, penflutazide, polythiazide, teclothiazide, trichlormethiazide, triflumethazide, and the like); ambuside, amiloride, aminometradine, azosemide, bemetizide, bumetanide, butazolamide, butizide, ethacrynic acid, canrenone, chloraminophenamide, chlorazanyl, chlormerodrin, chlorthalidone, clofenamide, clopamide, clorexolone, disulfamide, ethacrynic acid, ethoxzolamide, etozolon, fenquizone, furosemide, mefruside, meralluride, mercaptopimerin sodium, mercumallylic acid, mersalyl, methazolamide, metolazone, muzolimine, pamabrom, paraflutizide, piretanide, protheobromine, quinethazone, scoparius, spironalactone, theobromine, ticrynafen, torsemide, triamterene, xipamide or potassium, and the like. Suitable diuretics are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable endothelin antagonists include, but are not limited to, bosentan, endothelin, sulfonamide endothelin antagonists, BQ-123, SQ 28608, and the like. Suitable endothelin antagonists are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable H₂ receptor antagonists include, but are not limited to, burimamide, cimetidine, ebrotidine, famotidine, nizatidine, roxatidine, rantidine, tiotidine, and the like. Suitable H₂ receptor antagonists are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995, Pgs. 901-915; the Merck Index on CD-ROM, 13th Edition; and in WO 00/28988 assigned to NitroMed Inc., the disclosures of which are incorporated herein by reference in their entirety.

Suitable neutral endopeptidase inhibitors include, but are not limited to, atrial natriuretic peptides, diazapins, azepinones, ecadotril, omapatrilat, sampatrilat, BMS 189,921, and the like. Neutral endopeptidase inhibitors are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable NSAIDs include, but are not limited to, acetaminophen, acetaminophen, aceclofenac, alminoprofen, amfenac, bendazac, benoxaprofen, bromfenac, bucloxic acid, butibufen, carprofen, cinmetacin, clopirac, diclofenac, etodolac, felbinac, fenclozic acid, fenbufen, fenoprofen, fentiazac, flunoxaprofen, flurbiprofen, ibufenac, ibuprofen, indomethacin, isofezolac, isoxepac, indoprofen, ketoprofen, lonazolac, loxoprofen, metiazinic acid, mofezolac, miroprofen, naproxen, oxaprozin, pirozolac, piroprofen, pranoprofen, protizinic acid, salicylamide, sulindac, suprofen, suxibuzone, tiaprofenic acid, tolmetin, xenbucin, ximoprofen, zaltoprofen, zomepirac, aspirin, acetaminophen, bumadizon, carprofenac, clidanac, diflunisal, enfenamic acid, fendosal, flufenamic acid, flunixin, gentisic acid, ketorolac, meclofenamic acid, mefenamic acid, mesalamine, prodrugs thereof, and the like. Suitable NSAIDs are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995, Pgs. 617-657; the Merck Index on CD-ROM, 13th Edition; and in U.S. Patent Nos. 6,057,347 and 6,297,260 assigned to NitroMed Inc., the disclosures of which are incorporated herein by reference in their entirety.

Suitable potassium channel blockers include but are not limited to, nicorandil, pinacidil, cromakalim (BRL 34915), aprikalim, bimakalim, emakalim, lemakalim, minoxidil, diazoxide, 9-chloro-7-(2-chlorophenyl)-5H-pyrimido(5,4-d)(2)-benzazepine, Ribit, CPG-11952, CGS-9896, ZD 6169, diazoxide, Bay X 9227, P1075, Bay X 9228, SDZ PCO 400, WAY-120,491, WAY-

120,129, Ro 31-6930, SR 44869, BRL 38226, S 0121, SR 46142A, CGP 42500, SR 44994, artilide fumarate, lorazepam, temazepam, rilmafazone, nimetazepam, midazolam, lormetazepam, loprazolam, ibutilide fumarate, haloxazolam, flunitrazepam, estazolam, doxefazepam, clonazepam, cinolazepam, brotizolam, and the like. Suitable potassium channel blockers are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable platelet reducing agents include but are not limited to, fibrinolytic agents such as for example, ancrod, anistreplase, bisobrin lactate, brinolase, Hageman factor (i.e. factor XII) fragments, molsidomine, plasminogen activators such as, for example, streptokinase, tissue plasminogen activators (TPA), urokinase, pro-Urokinase, recombinant TPA, plasmin, plasminogen, and the like; anti-coagulant agents including but are not limited to, inhibitors of factor Xa, factor TFPI, factor VIIa, factor IXc, factor Va, factor VIIIa, inhibitors of other coagulation factors, and the like; vitamin K antagonists, such as, for example, coumarin, coumarin derivatives (e.g., warfarin sodium); glycosaminoglycans such as, for example, heparins both in unfractionated form and in low molecular weight form; ardeparin sodium, bivalirudin, bromindione, coumarin, dalteparin sodium, danaparoid sodium; dazoxiben hydrochloride, desirudin, dicumarol, efegatran sulfate, enoxaparin sodium, ifetroban, ifetroban sodium, lyapolate sodium, nafamostat mesylate, phenprocoumon, sulfatide, tinzaparin sodium, retaplast; trifluorethyl, warfarin, dextrans and the like; acadesine, anipamil, argatroban, aspirin, clopidogrel, diadenosine 5',5'''-P₁,P₄-tetrphosphate (Ap₄A) analogs, difibrotide, dilazep dihydrochloride, dipyridamole, dopamine, 3-methoxytyramine, glucagon, glycoprotein IIb/IIIa antagonists, such as, for example, Ro-43-8857, L-700,462, iloprost, isocarbacyclin methyl ester, itazigrel, ketanserin, BM-13.177, lamifiban, lifarizine, molsidomine, nifedipine, oxagrelate, prostaglandins, platelet activating factor antagonists such as, for example, lexipafant, prostacyclins, pyrazines, pyridinol carbamate, ReoPro (i.e., abciximab), sulfinpyrazone, synthetic compounds BN-50727, BN-52021, CV-4151, E-5510, FK-409, GU-7, KB-2796, KBT-3022, KC-404, KF-4939, OP-41483, TRK-100, TA-3090, TFC-612, ZK-36374, 2,4,5,7-tetrathiaoctane, 2,4,5,7-tetrathiaoctane 2,2-dioxide, 2,4,5-trithiahexane, theophyllin pentoxifyllin, thromboxane and thromboxane synthetase inhibitors such as, for example, picotamide, sulotroban, ticlopidine, tirofiban, trapidil, ticlopidine, trifluorethyl, trilineolein, 3-substituted 5,6-bis(4-methoxyphenyl)-

1,2,4-triazines; antibodies to glycoprotein IIb/IIIa; anti-serotonin drugs, such as, for example, clopidogrel; sulfinpyrazone and the like; aspirin; dipyridamole; clofibrate; pyridinol carbamate; glucagon, caffeine; theophyllin pentoxifyllin; ticlopidine, and the like.

Suitable proton pump inhibitors include, but are not limited to, omeprazole, lansoprazole, pantoprazole, rabeprazole, leminoprazole, timoprazole, tenatoprazole, disulprazole, esomeprazole, 2-(2-benzimidazolyl)-pyridine, tricyclic imidazole, thienopyridine benzimidazole, fluoroalkoxy substituted benzimidazole, dialkoxyl benzimidazole, N-substituted 2-(pyridylalkenesulfinyl) benzimidazole, cycloheptenepyrindine, 5-pyrrolyl-2-pyridylmethylsulfinyl benzimidazole, alkylsulfinyl benzimidazole, fluoro-pyridylmethylsulfinyl benzimidazole, imidazo[4,5-b]pyridine, RO 18-5362, IY 81149, 4-amino-3-carbonyl quinoline, 4-amino-3-acylnaphthyridine, 4-aminoquinoline, 4-amino-3-acylquinoline, 3-butyryl-4-(2-methylphenylamino)-8-(2-hydroxyethoxy)quinoline, quinazoline, tetrahydroisoquinolin-2-yl pyrimidine, YH 1885, 3-substituted 1,2,4-thiadiazolo[4,5-a] benzimidazole, 3-substituted imidazo[1,2-d]-thiadiazole, 2-sulfinylnicotinamide, pyridylsulfinylbenzimidazole, pyridylsulfinyl thieno imidazole, theinoimidazole-toluidine, 4,5-dihydrooxazole, thienoimidazole-toluidine, Hoe-731, imidazo[1,2-a]pyridine, pyrrolo[2,3-b]pyridine, and the like.

Suitable proton pump inhibitors are described more fully in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill, 1995; the Merck Index on CD-ROM, 13th Edition; and in WO 00/50037 assigned to NitroMed Inc., the disclosures of which are incorporated herein by reference in their entirety.

Suitable renin inhibitors include, but are not limited to, aldosterone, enalkrein, medullipin, tonin, RO 42-5892, A 65317, CP 80794, ES 1005, ES 8891, SQ 34017, urea derivatives of peptides, amino acids connected by nonpeptide bonds, di- and tri-peptide derivatives, amino acids and derivatives thereof, diol sulfonamides and sulfinyls, modified peptides, peptidyl beta-aminoacyl aminodiol carbamates, monoclonal antibodies to rennin, and the like. Suitable renin inhibitors are described more fully in U.S. Patent Nos. 5,116,835, 5,114,937, 5,106,835, 5,104,869, 5,095,119, 5,098,924, 5,095,006, 5,089,471, 5,075,451, 5,066,643, 5,063,208, 4,845,079, 5,055,466, 4,980,283, 4,885,292, 4,780,401, 5,071,837, 5,064,965, 5,063,207, 5,036,054, 5,036,053, 5,034,512, and 4,894,437, the disclosures of each of which are incorporated herein by reference in their entirety; and in the literature, such as in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (9th Edition), McGraw-Hill,

1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Suitable COX-2 inhibitors include, but are not limited to, NS-386, nimesulide, flosulide, celecoxib, rofecoxib, COX-189, etoracoxib, Bextra, Dynastat, Arcoxia, SC-57666, DuP 697, SC-58125, SC-58635, and the like, and mixtures of two or more thereof. Suitable COX-2 inhibitors are in U.S. Patent Nos. 5,344,991, 5,380,738, 5,393,790, 5,409,944, 5,434,178, 5,436,265, 5,466,823, 5,474,995, 5,510,368, 5,536,752, 5,550,142, 5,552,422, 5,604,253, 5,604,260, and 5,639,780 and in WO 94/03387, WO 94/15723, WO 94/20480, WO 94/26731, WO 94/27980, WO 95/00501, WO 95/15316, WO 96/03387, WO 96/03388, WO 96/06840, WO 96/21667, WO 96/31509, WO 96/36623, WO 97/14691, WO 97/16435, WO 01/45703 and WO 01/87343, the disclosures of each of which are incorporated herein by reference in their entirety; and in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, 1995; and the Merck Index on CD-ROM, Thirteenth Edition; and on STN Express, file phar and file registry.

Another embodiment of the invention provides methods for preventing and/or treating cardiovascular diseases by administering to the patient in need thereof a therapeutically effective amount of the compounds and/or compositions described herein. For example, the patient can be administered a therapeutically effective amount of at least one nitrosated and/or nitrosylated cardiovascular compound. In another embodiment, the patient can be administered a therapeutically effective amount of at least one cardiovascular compound, that is optionally nitrosated and/or nitrosylated, and at least one compound that donates, transfers or releases nitric oxide, or elevates levels of endogenous EDRF or nitric oxide, or is a substrate for nitric oxide synthase. In yet another embodiment, the patient can be administered a therapeutically effective amount of at least one cardiovascular compound, that is optionally nitrosated and/or nitrosylated, and, at least one therapeutic agent, including but not limited to, such as, for example, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H₂ receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin

inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and, optionally, at least one compound that donates, transfers or releases nitric oxide, or elevates levels of endogenous EDRF or nitric oxide, or is a substrate for nitric oxide synthase. The compounds can be administered separately or in the form of a composition.

5 Yet another embodiment of the invention provides methods for treating diseases resulting from oxidative stress by administering to the patient in need thereof a therapeutically effective amount of the compounds and/or compositions described herein. For example, the patient can be administered a therapeutically effective amount of at least one nitrosated and/or nitrosylated cardiovascular compound. In another embodiment, the patient can be administered a
10 therapeutically effective amount of at least one cardiovascular compound, that is optionally nitrosated and/or nitrosylated, and at least one compound that donates, transfers or releases nitric oxide, or elevates levels of endogenous EDRF or nitric oxide, or is a substrate for nitric oxide synthase. In yet another embodiment, the patient can be administered a therapeutically effective amount of at least one cardiovascular compound, that is optionally nitrosated and/or nitrosylated,
15 and, at least one therapeutic agent, including but not limited to, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H₂ receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal
20 antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and, optionally, at least one nitric oxide donor. The compounds can be administered separately or in the form of a composition.

Yet another embodiment of the invention provides methods for treating endothelial
25 dysfunctions by administering to the patient in need thereof a therapeutically effective amount of the compounds and/or compositions described herein. For example, the patient can be administered a therapeutically effective amount of at least one nitrosated and/or nitrosylated cardiovascular compound. In another embodiment, the patient can be administered a therapeutically effective amount of at least one cardiovascular compound, that is optionally
30 nitrosated and/or nitrosylated, and at least one compound that donates, transfers or releases nitric oxide, or elevates levels of endogenous EDRF or nitric oxide, or is a substrate for nitric oxide

synthase. In yet another embodiment, the patient can be administered a therapeutically effective amount of at least one cardiovascular compound, that is optionally nitrosated and/or nitrosylated, and, at least one therapeutic agent, including but not limited to, aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H₂ receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and, optionally, at least one nitric oxide donor. The compounds can be administered separately or in the form of a composition.

When administered separately, the cardiovascular compound, that is optionally nitrosated and/or nitrosylated, nitric oxide donor and/or therapeutic agent can be administered about the same time as part of the overall treatment regimen, i.e., as a combination therapy. "About the same time" includes administering the cardiovascular compound, that is optionally nitrosated and/or nitrosylated, simultaneously, sequentially, at the same time, at different times on the same day, or on different days, as long as they are administered as part of an overall treatment regimen, i.e., combination therapy or a therapeutic cocktail.

When administered in vivo, the compounds and compositions of the invention can be administered in combination with pharmaceutically acceptable carriers and in dosages described herein. When the compounds and compositions of the invention are administered as a combination of at least one cardiovascular compound and/or at least one nitrosated and/or nitrosylated cardiovascular compound and/or at least one nitric oxide donor and/or therapeutic agent, they can also be used in combination with one or more additional compounds which are known to be effective against the specific disease state targeted for treatment. The nitric oxide donors, therapeutic agents and/or other additional compounds can be administered simultaneously with, subsequently to, or prior to administration of the nitrosated and/or nitrosylated cardiovascular compound.

The compounds and compositions of the invention can be administered by any available and effective delivery system including, but not limited to, orally, buccally, parenterally, by inhalation spray, by topical application, by injection, transdermally, or rectally (e.g., by the use of

suppositories) in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles, as desired. Parenteral includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Transdermal compound administration, which is known to one skilled in the art, involves the delivery of pharmaceutical compounds via percutaneous passage of the compound into the systemic circulation of the patient. Topical administration can also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. Other components can be incorporated into the transdermal patches as well. For example, compositions and/or transdermal patches can be formulated with one or more preservatives or bacteriostatic agents including, but not limited to, methyl hydroxybenzoate, propyl hydroxybenzoate, chlorocresol, benzalkonium chloride, and the like. Dosage forms for topical administration of the compounds and compositions can include creams, sprays, lotions, gels, ointments, eye drops, nose drops, ear drops, and the like. In such dosage forms, the compositions of the invention can be mixed to form white, smooth, homogeneous, opaque cream or lotion with, for example, benzyl alcohol 1% or 2% (wt/wt) as a preservative, emulsifying wax, glycerin, isopropyl palmitate, lactic acid, purified water and sorbitol solution. In addition, the compositions can contain polyethylene glycol 400. They can be mixed to form ointments with, for example, benzyl alcohol 2% (wt/wt) as preservative, white petrolatum, emulsifying wax, and tenox II (butylated hydroxyanisole, propyl gallate, citric acid, propylene glycol). Woven pads or rolls of bandaging material, e.g., gauze, can be impregnated with the compositions in solution, lotion, cream, ointment or other such form can also be used for topical application. The compositions can also be applied topically using a transdermal system, such as one of an acrylic-based polymer adhesive with a resinous crosslinking agent impregnated with the composition and laminated to an impermeable backing.

The compositions can also be applied topically using a transdermal system, such as one of an acrylic-based polymer adhesive with a resinous crosslinking agent impregnated with the composition and laminated to an impermeable backing. In a particular embodiment, the compositions of the invention are administered as a transdermal patch, more particularly as a sustained-release transdermal patch. The transdermal patches of the invention can include any conventional form such as, for example, adhesive matrix, polymeric matrix, reservoir patch, matrix or monolithic-type laminated structure, and are generally comprised of one or more

backing layers, adhesives, penetration enhancers, an optional rate controlling membrane and a release liner which is removed to expose the adhesives prior to application. Polymeric matrix patches also comprise a polymeric-matrix forming material. Suitable transdermal patches are described in more detail in, for example, U. S. Patent Nos. 5,262,165, 5,948,433, 6,010,715 and 6,071,531, the disclosure of each of which are incorporated herein in their entirety.

Solid dosage forms for oral administration can include capsules, sustained-release capsules, tablets, sustained release tablets, chewable tablets, sublingual tablets, effervescent tablets, pills, powders, granules and gels. In such solid dosage forms, the active compounds can be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms can also comprise, as in normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, effervescent tablets, and pills, the dosage forms can also comprise buffering agents. Soft gelatin capsules can be prepared to contain a mixture of the active compounds or compositions of the invention and vegetable oil. Hard gelatin capsules can contain granules of the active compound in combination with a solid, pulverulent carrier such as lactose, saccharose, sorbitol, mannitol, potato starch, corn starch, amylopectin, cellulose derivatives of gelatin. Tablets and pills can be prepared with enteric coatings.

Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions can also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

Suppositories for vaginal or rectal administration of the compounds and compositions of the invention, such as for treating pediatric fever and the like, can be prepared by mixing the compounds or compositions with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at room temperature but liquid at rectal temperature, such that they will melt in the rectum and release the drug.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable dispersing agents, wetting agents and/or suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be used are

water, Ringer's solution, and isotonic sodium chloride solution. Sterile fixed oils are also conventionally used as a solvent or suspending medium.

The compositions of this invention can further include conventional excipients, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral application which do not deleteriously react with the active compounds. Suitable pharmaceutically acceptable carriers include, for example, water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, surfactants, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, and the like. The pharmaceutical preparations can be sterilized and if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds. For parenteral application, particularly suitable vehicles consist of solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants. Aqueous suspensions may contain substances which increase the viscosity of the suspension and include, for example, sodium carboxymethyl cellulose, sorbitol and/or dextran. Optionally, the suspension may also contain stabilizers.

The composition, if desired, can also contain minor amounts of wetting agents, emulsifying agents and/or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like.

Various delivery systems are known and can be used to administer the compounds or compositions of the invention, including, for example, encapsulation in liposomes, microbubbles, emulsions, microparticles, microcapsules and the like. The required dosage can be administered as a single unit or in a sustained release form.

The bioavailability of the compositions can be enhanced by micronization of the formulations using conventional techniques such as grinding, milling, spray drying and the like in the presence of suitable excipients or agents such as phospholipids or surfactants.

4
-
Sustained release dosage forms of the invention may comprise microparticles and/or nanoparticles having a therapeutic agent dispersed therein or may comprise the therapeutic agent in pure, preferably crystalline, solid form. For sustained release administration, microparticle dosage forms comprising pure, preferably crystalline, therapeutic agents are preferred. The therapeutic dosage forms of this aspect of the invention may be of any configuration suitable for sustained release.

5
Nanoparticle sustained release therapeutic dosage forms are preferably biodegradable and, optionally, bind to the vascular smooth muscle cells and enter those cells, primarily by endocytosis. The biodegradation of the nanoparticles occurs over time (e.g., 30 to 120 days; or 10 to 21 days) in prelysosomal vesicles and lysosomes. Preferred larger microparticle therapeutic dosage forms of the invention release the therapeutic agents for subsequent target cell uptake with only a few of the smaller microparticles entering the cell by phagocytosis. A practitioner in the art will appreciate that the precise mechanism by which a target cell assimilates and metabolizes a dosage form of the invention depends on the morphology, physiology and metabolic processes of those cells. The size of the particle sustained release therapeutic dosage forms is also important with respect to the mode of cellular assimilation. For example, the smaller nanoparticles can flow with the interstitial fluid between cells and penetrate the infused tissue. The larger microparticles tend to be more easily trapped interstitially in the infused primary tissue, and thus are useful to deliver anti-proliferative therapeutic agents.

10
15
20
Particular sustained release dosage forms of the invention comprise biodegradable microparticles or nanoparticles. More particularly, biodegradable microparticles or nanoparticles are formed of a polymer containing matrix that biodegrades by random, nonenzymatic, hydrolytic scissioning to release therapeutic agent, thereby forming pores within the particulate structure.

25
In a particular embodiment, the compositions of the invention are orally administered as a sustained release tablet or a sustained release capsule. For example, the sustained release formulations can comprise a therapeutically effective amount of at least one nitrosated and/or nitrosylated cardiovascular compound or a pharmaceutically acceptable salt thereof, and, optionally at least one nitric oxide donor, or the sustained release formulations can comprise a therapeutically effective amount of at least one nitrosated and/or nitrosylated cardiovascular

30

compound or a pharmaceutically acceptable salt thereof, and at least one nitric oxide donor, and, optionally at least one therapeutic agent

The compounds and compositions of the invention can be formulated as pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include, for example, alkali metal salts and addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids include, but are not limited to, hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid and the like. Appropriate organic acids include, but are not limited to, aliphatic, cycloaliphatic, aromatic, heterocyclic, carboxylic and sulfonic classes of organic acids, such as, for example, formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, algenic, β -hydroxybutyric, cyclohexylaminosulfonic, galactaric and galacturonic acid and the like. Suitable pharmaceutically-acceptable base addition salts include, but are not limited to, metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from primary, secondary and tertiary amines, cyclic amines, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine and the like. All of these salts may be prepared by conventional means from the corresponding compound by reacting, for example, the appropriate acid or base with the compound.

While individual needs may vary, determination of optimal ranges for effective amounts of the compounds and/or compositions is within the skill of the art. Generally, the dosage required to provide an effective amount of the compounds and compositions, which can be adjusted by one of ordinary skill in the art, will vary depending on the age, health, physical condition, sex, diet, weight, extent of the dysfunction of the recipient, frequency of treatment and the nature and scope of the dysfunction or disease, medical condition of the patient, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of the particular compound used, whether a drug delivery system is used, and whether the compound is administered as part of a drug combination.

The amount of a given nitrosated and/or nitrosylated cardiovascular compound of the invention that will be effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques, including reference to Goodman and Gilman, supra; The Physician's Desk Reference, Medical Economics Company, Inc., Oradell, N.J., 1995; and Drug Facts and Comparisons, Inc., St. Louis, MO, 1993. The precise dose to be used in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided by the physician and the patient's circumstances.

The invention also provides pharmaceutical kits comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compounds and/or compositions of the invention, including, at least, one or more of the novel cardiovascular compound, that is optionally nitrosated and/or nitrosylated, and one or more of the NO donors described herein. Associated with such kits can be additional therapeutic agents or compositions (e.g., aldosterone antagonists, alpha-adrenergic receptor antagonists, antidiabetic compounds, anti-hyperlipidemic drugs, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, antioxidants, antithrombotic and vasodilator drugs, beta-adrenergic blockers, calcium channel blockers, diuretics, endothelin antagonists, H₂ receptor antagonists, neutral endopeptidase inhibitors, nonsteroidal antiinflammatory compounds (NSAIDs), potassium channel blockers, platelet reducing agents, proton pump inhibitors, renin inhibitors, selective cyclooxygenase-2 (COX-2) inhibitors, and the like, and mixtures of two or more thereof), devices for administering the compositions, and notices in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products which reflects approval by the agency of manufacture, use or sale for humans.

The disclosure of each patent, patent application and publication cited or described in the present specification is hereby incorporated by reference herein in its entirety.

Although the invention has been set forth in detail, one skilled in the art will appreciate that numerous changes and modifications can be made to the invention, and that such changes and modifications can be made without departing from the spirit and scope of the invention.

CLAIMS

What is claimed is:

1. A nitrosated and/or nitrosylated aldosterone antagonist, a nitrosated and/or nitrosylated angiotensin II antagonist, a nitrosated and/or nitrosylated hydralazine compound, a nitrosated and/or nitrosylated calcium channel blocker, a nitrosated and/or nitrosylated diuretic, a nitrosated and/or nitrosylated endothelin antagonist, a nitrosated and/or nitrosylated neutral endopeptidase inhibitor, a nitrosated and/or nitrosylated renin inhibitor or a pharmaceutically acceptable salt thereof, that must contain at least at least one nitrite, nitrate, thionitrate or thionitrite group.

10

ABSTRACT OF THE DISCLOSURE

The invention describes novel nitrosated and/or nitrosylated cardiovascular compounds and pharmaceutically acceptable salts thereof, and novel compositions comprising at least one nitrosated and/or nitrosylated cardiovascular compound, and, optionally, at least one compound that donates, transfers or releases nitric oxide, stimulates endogenous synthesis of nitric oxide, elevates endogenous levels of endothelium-derived relaxing factor or is a substrate for nitric oxide synthase, and/or at least one therapeutic agent. The invention also provides novel compositions comprising at least one cardiovascular compound, and at least one nitric oxide donor and/or at least one therapeutic agent. The invention also provides methods for treating cardiovascular diseases, methods for treating diseases resulting from oxidative stress and methods for treating endothelial dysfunctions. The nitrosated and/or nitrosylated cardiovascular compounds are preferably nitrosated and/or nitrosylated aldosterone antagonists, nitrosated and/or nitrosylated angiotensin II antagonists, nitrosated and/or nitrosylated hydralazine compounds, nitrosated and/or nitrosylated calcium channel blockers, nitrosated and/or nitrosylated diuretics, nitrosated and/or nitrosylated endothelin antagonists, nitrosated and/or nitrosylated neutral endopeptidase inhibitors and nitrosated and/or nitrosylated renin inhibitors.

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/026910

International filing date: 20 August 2004 (20.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/498,309
Filing date: 28 August 2003 (28.08.2003)

Date of receipt at the International Bureau: 15 December 2004 (15.12.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse